

How to use DAIVOBET® Gel



1

Shake the bottle before use and remove the cap. Comb the hair first to remove any loose scales (it may help to part your hair before you apply the gel).



4

In order to achieve optimal effects, it is recommended that the hair is not washed immediately after application of Daivobet® gel. Let the gel remain on the scalp during the night or during the day.




2

Apply a drop of Daivobet® gel to the fingertip. Usually an amount between 1g and 4g per day is sufficient treatment for the scalp. Tilt your head to make sure the gel does not run onto your face.



5

When washing hair, apply a mild shampoo to **DRY HAIR**, especially to those areas where the gel was applied.



3

Apply the gel to the areas of the scalp affected by psoriasis and gently rub in with your fingertips. Wash your hands thoroughly after applying the gel. This will avoid spreading the gel to other parts of the body (especially the face, mouth and eyes).



6

Leave the shampoo on the scalp for a couple of minutes before washing your hair as usual.

DAIVOBET® Gel and Ointment are fully funded.

PRESENTATION: DAIVOBET® Ointment and Gel contains 50 mcg/g calcipotriol and 500 mcg/g betamethasone (as dipropionate); available in 30g tubes and bottles, respectively. **INDICATION:** Treatment of plaque-type psoriasis vulgaris and scalp psoriasis in adult patients. **DOSAGE AND ADMINISTRATION:** Apply topically once daily. The maximum daily dose should not exceed 15 grams. **CONTRAINDICATIONS:** Allergy to any constituent of the product; disorders of calcium metabolism; viral, fungal or bacterial skin infections; parasitic infections; tuberculosis or syphilis; perioral dermatitis; acne vulgaris; atrophic skin; striae atrophicae; fragility of skin veins; ichthyosis; acne rosacea; ulceration; wounds; perianal and genital pruritus; guttate, erythrodermic, exfoliative and pustular psoriasis; severe renal insufficiency or hepatic disorders. **WARNINGS AND PRECAUTIONS:** Face; mouth; eyes; genitals; more than 30% of the body surface area; serum calcium and renal function; concurrent treatment with other steroids; occlusive dressings; large areas of damaged skin; mucous membranes; skin folds; sunlight and UV; long-term use; pregnancy; lactation; breast area. **ADVERSE EFFECTS:** Pruritus; rash scaly; ecchymosis; headache; nasopharyngitis; burning and stinging sensation of the skin; skin pain or irritation; dermatitis; erythema; acne; exacerbation of psoriasis; folliculitis; transient photosensitivity; depigmentation; dry skin; eczema; hypercalcaemia or hypercalciuria; skin atrophy; telangiectasia; striae; hypertrichosis; colloid milia; adrenocortical suppression; cataract; infections; increase in intra-ocular pressure; HPA suppression. **INTERACTIONS:** Calcium or vitamin D supplements. **MEDICINE CLASSIFICATION:** Prescription Medicine. Please review the full data sheets before prescribing DAIVOBET®. Full data sheets are available from CSL Biotherapies (NZ) Ltd, PO Box 62 590, Greenlane, Auckland 1546, www.csllbiotherapies.co.nz, or the Medsafe website www.medsafe.govt.nz

Based on the DAIVOBET® Ointment and Gel data sheets dated December 2011. DAIVOBET® is a registered trademark of Leo Pharmaceutical Products Ltd A/S. DBET-022-2/12. DA1111PG



Daivobet®
calcipotriol /
betamethasone dipropionate

CSL Biotherapies